Executive Summary:

Proven Surgical Nutrition Protocol with IMPACTAdvanced Recovery® Drink



Concept

• A surgical nutrition protocol incorporating **IMPACT Advanced Recovery® Drink** provides evidence based immune support for major elective surgery patients to reduce the risk of costly postoperative infectious complications, length of stay (LOS) and readmissions. ¹⁻³

Challenges Addressed by IMPACT® Surgical Nutrition

- The Hospital Acquired Condition (HAC) Reduction Program looks at reducing 6 key measures:4
- 1 Patient safety indicator (PSI) 90 composite measure (including wound dehiscence rate and pressure injury)
- **2 -** Central line associated bloodstream infections (CLABSI)
- **3 -** Catheter-associated urinary tract infections (CAUTIs)
- 4 Surgical site infections (SSIs) specifically in colonic surgeries and abdominal hysterectomies
- **5 -** Methicillin-resistant *Staphlococcus aureus* (MRSA) bacteremia
- **6 -** Clostridium difficile infection (CDI)
- A large retrospective analysis of major elective surgery patients showed a preoperative kit, including IMPACT Advanced Recovery® Drink, helped to reduce SSI (p<0.044), CAUTI (p<0.007), CDI (p<0.016), and PSI (p<0.001) in the intervention vs. pre-intervention group.⁵
- The cost of a single SSI can be more than \$20,000,6 therefore a decrease in SSIs can have a significant effect on helping health systems achieve the Triple Aim (Better Health, Better Care, Lower Costs).⁷

Evidence Supporting IMPACT® Surgical Nutrition

- The blend of immunonutrients in **IMPACT Advanced Recovery® Drink is supported by 40+ randomized controlled trials and multiple quality improvement initiatives** across multiple surgery types: GI, cardiac, head and neck cancer, bladder cancer, gyn-oncology, orthopedic, thoracic cancer and ventral hernia repair.^{5,8-16}
- IMPACT Advanced Recovery® Drink contains immunonutrients recommended perioperatively by the Society of Critical Care Medicine (SCCM) and the American Society of Parenteral And Enteral Nutrition (A.S.P.E.N), the A.S.P.E.N. Guidelines on Nutritional Support Therapy During Adult Anti-Cancer Treatment, the Guidelines on Enteral Nutrition: Surgery (ESPEN), the American Society of Enhanced Recovery (ASER) Implementation Guide, and the Perioperative Quality Initiative (POQI) Joint Consensus Statement.^{17-21*}
- A 50-58% reduction in hospital readmission of colorectal surgery patients was found with the use of the Strong for Surgery** (S4S) Nutrition Checklist and incorporation of preoperative IMPACT Advanced Recovery®.2, 22
 - Mean total costs were \$2,500 less at discharge for colorectal surgery patients who were S4S checklisted and supplemented with IMPACT Advanced Recovery[®].²

What Makes IMPACT® Surgical Nutrition Unique

- The synergistic blend of arginine, omega-3 fatty acids from fish oil, and dietary nucleotides supports the unique nutritional needs of the major elective surgery patient.²³⁻²⁵
- Only this unique blend of ingredients is proven to:
 - Reduce the risk of postoperative infectious complications by 51% (p<0.00001) and hospital length of stay by 15% (avg. 2–3 days) (p<0.00001) after major elective surgery.¹
 - Reduce the risk of postoperative complications such as anastomotic leak, UTI, nosocomial pneumonia, pharyngeal fistula, wound complications and surgical site infections (SSIs).^{9, 11, 26-28}

Benefits of Using IMPACT® Surgical Nutrition as part of an Enhanced Recovery Protocol (ERP)

- ERPs that include IMPACT Advanced Recovery® have shown reductions in LOS, readmissions, hospital charges and infection rates. 12, 15, 16, 29
- GI surgery patients provided IMPACT Advanced Recovery® Drink preoperatively as part of an ERP, and at no charge, had a significantly reduced LOS when compared to ERP patients purchasing the product out-of-pocket.³⁰

Implementation

- Perioperative use of IMPACT Advanced Recovery® Drink is recommended for patients undergoing major elective surgery. Recommend three cartons per day or one UltraPak® per day*** for five days prior to surgery and for at least five days following surgery.^{9, 27, 29}
- A Nestlé Representative supports implementation of the perioperative protocol by providing program awareness, staff education and compliance support.
- *This statement does not constitute an endorsement of IMPACT Advanced Recovery® Drink or any other Nestlé Health Science product by SCCM, A.S.P.E.N., ESPEN or ASER.
- ** The American College of Surgeons and Strong for Surgery do not endorse IMPACT® formulas or any other product.
- Strong for Surgery is a Quality Program of the American College of Surgeons.

^{***} Initiate IMPACT® Peptide 1.5 or IMPACT® tube feeding formula and advance to ≥1,000 calories/day to meet nutritional needs. Also available in open system.

References

- **1.** Drover JW et al. Perioperative use of arginine-supplemented diets: A systematic review of the evidence. J Am Coll Surg 2011; 212(3): 385-399.
- 2. Banerjee S et al. Effects of Arginine-based Immunonutrition on Inpatient Total Costs and Hospitalization Outcomes for Patients Undergoing Colorectal Surgery. *Nutr* 2017;42:106-13.
- **3.** Mauskopf JA et al. Immunonutrition for patients undergoing elective surgery for gastrointestinal cancer: Impact on hospital costs. *WJSO* 2012;10:136.
- 4. https://www.cms.gov/Medicare/Medicare-Fee-for-Service-Payment/AcuteInpatientPPS/Downloads/FY2018-HAC-Reduction-Program-Fact-Sheet.pdf
- 5. Kelley KE et al. Impact of a Novel Preoperative Patient-Centered Surgical Wellness Program. Ann Surg 2018;268(4): 650-56.
- **6.** de Lissovoy G et al. Surgical site infection: incidence and impact on hospital utilization and treatment costs. *Am J Infect Control* 2009; 37:387-397.
- 7. Triple Aim for Populations. www.ihi.org/Topics/TripleAim. Accessed May 25, 2016.
- **8.** Tepaske R et al. Glycine does not add to the beneficial effects of perioperative oral immune-enhancing nutrition supplements in high-risk cardiac surgery patients. *JPEN* 2007;31(3):173-180.
- **9.** Rowan NR et al. Utility of a Perioperative Nutritional Intervention on Postoperative Outcomes in High-Risk Head and Neck Cancer Patients. *Oral Onc* 2016; 54:42-46.
- **10.** Bertrand BJ et al. Impact of Preoperative Immunonutrition on Morbidity Following Cystectomy for Bladder Cancer: A Case-Control Pilot Study. *World J Urol* 2014; 32:233-237.
- **11.** Chapman JS et al. Post-operative Enteral Immunonutrition for Gynecologic Oncology Patients Undergoing Laparotomy Decreases Wound Complications. *Gynecologic Oncology* 2015;137:523-528.
- **12.** Alito Aprelino M and de Aguilar-Nascimento JE. Multimodal Perioperative Care plus Immunonutrition versus Traditional Care in Total Hip Arthroplasty: A Randomized Pilot Study. *Nutrition Journal* 2016;15:1-7.
- 13. Heynen K. Pre-op Nutrition: Making an "IMPACT". Presented at the American Society of Enhanced Recovery (ASER), May 15, 2018.
- **14.** Kaya SO et al. Is preoperative protein-rich nutrition effective on postoperative outcome in non-small cell lung cancer surgery? A prospective randomized study *J Cardiothor Surg* 2016;11:14.
- **15.** Majumder A et al. Benefits of Multimodal Enhanced Recovery Pathway in Patients Undergoing Open Ventral Hernia Repair. J Am Coll Surg 2016;222:1106-1115.
- **16.** Robinson LA et al. Preop Nutrition-Enhanced Recovery After Surgery (N-ERAS©) Protocol for Thoracic Cancer Resections Decreases Hospital Days, Complications and Charges. Presented at IASLC Sept 2018.
- **17.** McClave SA et al. Guidelines for the provision and assessment of nutrition support therapy in the adult critically ill patient: SCCM and A.S.P.E.N. *JPEN* 2009; 33(3): 277-316.
- **18.** August DA and Huhmann MB. A.S.P.E.N. Clinical Guidelines: Nutrition support therapy during adult anticancer treatment and in hematopoietic cell transplantation. *JPEN* 2009; 33(5): 472-500.
- 19. Weimann A et al. ESPEN guideline: Clinical nutrition in surgery. ESPEN 2017;36:623-650.
- 20. ASER. Enhanced Recovery Implementation Guide. Sept 2016. http://aserhq.org/membership-education/non-member-items/implementation-guide-preview/
- 21. Wischmeyer PE et al. American Society for Enhanced Recovery and Perioperative Quality Initiative Joint Consensus Statement on Nutrition Screening and Therapy with a Surgical Enhanced Recovery Pathway. Anesth Analg 2018;126(6):1883-95.
- 22. Strong for Surgery. American College of Surgeons. May 2017. http://www.facs.org/quality-programs/strong-for-surgery
- **23.** Zhu X et al. The central role of arginine catabolism on T-cell dysfunction and increased susceptibility to infection after physical injury. *Ann Surg* 2014;259;171-178.
- 24. Santora R and Kozar RA. Molecular Mechanisms of Pharmaconutrients. J Surg Res 2010;161:288-294.
- **25.** Bansal V et al. Interactions between Fatty Acids and Arginine Metabolism: Implications for the Design of Immune-Enhancing Diets. *JPEN* 2005: 29:S75.
- **26.** Horie H et al. Favorable effects of preoperative enteral immunonutrition on SSI infection in patients with colorectal cancer without malnutrition. *Surg Today* 2006;36:1063-1068.
- **27.** Waitzberg DL, Saito H et al. Postsurgical infections are reduced with specialized nutrition support. *World J Surg* 2006; 30:1592-1604.
- **28.** Farreras N et al. Effect of Early Postoperative Enteral Immunonutrition on Wound Healing in Patients Undergoing Surgery for Gastric Cancer. *Clin Nutr* 2005; 24:55-65.
- **29.** Hamilton-Reeves JM et al. Effects of Immunonutrition for Cystectomy on Immune Response and Infection Rates: A Pilot Randomized Controlled Clinical Trial. *Euro Urol* 2016;69(3):389-392.
- **30.** Cavanaugh E et al. Enhanced Recovery After Surgery: Does the Provision of an Immune-Modulating Protein Supplement Improve Post-Operative Outcomes? Presented at ERAS-USA Nov 2018.



