

Addressing Sarcopenia: Optimizing Protein Intake with Aging

References

Arentson-Lantz EJ, Galvan E, Ellison J, et al. Improving Dietary Protein Quality Reduces the Negative Effects of Physical Inactivity on Body Composition and Muscle Function. *The Journals of Gerontology: Series A*. 2019;74(10):1605-1611.

Bauer J, Biolo G, Cederholm T, et al. Evidence-based recommendations for optimal dietary protein intake in older people: a position paper from the PROT-AGE Study Group. *J Am Med Dir Assoc*. 2013;14(8):542-559.

Cruz-Jentoft AJ, Bahat G, Bauer J, et al. Sarcopenia: revised European consensus on definition and diagnosis. *Age Ageing*. 2019;48(1):16–31.

Deutz NE, Bauer JM, Barazzoni R, et al. Protein intake and exercise for optimal muscle function with aging: recommendations from the ESPEN Expert Group. *Clin Nutr*. 2014;33(6):929-936.

English KL, Paddon-Jones D. Protecting muscle mass and function in older adults during bed rest. *Curr Opin Clin Nutr Metab Care*. 2010;13(1):34-39.

English KL, Mettler JA, Ellison JB, Mamerow MM, Arentson-Lantz E, Pattarini JM, Ploutz-Snyder R, Sheffield-Moore M, Paddon-Jones D. Leucine partially protects muscle mass and function during bed rest in middle-aged adults. *Am J Clin Nutr*. 2016;103(2):465-73.

Fisher GG, Franks MM, Plassman BL, et al. Caring for individuals with dementia and cognitive impairment, not dementia: findings from the aging, demographics, and memory study. *J Am Geriatr Soc*. 2011;59(3):488-494.

Gross LS, Li L, Ford ES, Liu S. Increased consumption of refined carbohydrates and the epidemic of type 2 diabetes in the United States: an ecologic assessment. *Am J Clin Nutr*. 2004;79(5):774–779.

Heyland DK, Stapleton R, Compher C. Should we prescribe more protein to critically ill patients? *Nutrients*. 2018;10(4):462. <https://doi.org/10.3390/nu10040462>

Johnson Stoklossa CA, Ghosh SS, Forhan M, et al. Poor physical function as a marker of sarcopenia in adults with class II/III obesity. *Curr Dev Nutr*. 2018;2:nzx008. DOI:10.3945/cdn.117.001743

Katsanos CS, Kobayashi H, Sheffield-Moore M, et al. Aging is associated with diminished accretion of muscle proteins after the ingestion of a small bolus of essential amino acids. *Am J Clin Nutr*. 2005;82(5):1065–1073.

Kortebein P, Ferrando A, Lombeida J, et al. Effect of 10 days of bed rest on skeletal muscle in healthy older adults. *JAMA*. 2007;297(16):1772-1774.

Addressing Sarcopenia: Optimizing Protein Intake with Aging

References Continued

Mamerow MM, Mettler JA, English KL, et al. Dietary protein distribution positively influences 24-h muscle protein synthesis in healthy adults. *J Nutr*. 2014;144(6):876–880.

Paddon-Jones D, Campbell WW, Jacques PF, et al. Protein and healthy aging. *AJCN*. 2015;101(6):1339S–1345S.

Paddon-Jones D, Rasmussen BB. Dietary protein recommendations and the prevention of sarcopenia. *Curr Opin Clin Nutr Metab Care*. 2009;12(1):86–90.

Paddon-Jones D, Sheffield-Moore M, Urban RJ, et al. Essential amino acid and carbohydrate supplementation ameliorates muscle protein loss in humans during 28 days bedrest. *J Clin Endocrinol Metab*. 2004;89(9):4351–4358.

Rodriguez NR, Di Marco NM, Langley S. American College of Sports Medicine position stand. Nutrition and athletic performance. American Dietetic Association; Dietitians of Canada; American College of Sports Medicine. *Med Sci Sports Exerc*. 2009 Mar;41(3):709–731.

Symons TB, Schutzler SE, Cocke TL, et al. Aging does not impair the anabolic response to a protein-rich meal. *Am J Clin Nutr*. 2007;86(2):451–456.

Symons TB, Sheffield-Moore M, Wolfe RR, Paddon-Jones D. A moderate serving of high-quality protein maximally stimulates skeletal muscle protein synthesis in young and elderly subjects. *J Am Diet Assoc*. 2009;109(9):1582–1586.

Symons TB, Sheffield-Moore TM, Mamerow MM, et al. The anabolic response to resistance exercise and a protein-rich meal is not diminished by age. *J Nutr Health and Aging*. 2011;15(5):376–381.

Traylor DA, Gorissen SHM, Phillips SM. Perspective: protein requirements and optimal intakes in aging: are we ready to recommend more than the recommended daily allowance? *Adv Nutr*. 2018;9(3):171–182.

van Vliet S, Burd NA, van Loon LJ. The skeletal muscle anabolic response to plant- versus animal-based protein consumption. *J Nutr*. 2015;145(9):1981–1991.

Zachwieja JJ, Smith SR, Lovejoy JC, et al. Testosterone administration preserves protein balance but not muscle strength during 28 days of bed rest. *J Clin Endocrinol Metab*. 1999;84(1):207–212.