


# Nutritional Management of Pressure Injury – Oral Nutritional Supplements



**USE UNDER MEDICAL SUPERVISION.**

This document is not intended as a substitute for clinical judgment or existing facility protocols.

Posthauer M et al. *Adv Skin Wound Care* 2015;28(4):175-88.

<http://www.npuap.org/resources/educational-and-clinical-resources/npuap-pressure-injury-stages/>

All trademarks are owned by Société des Produits Nestlé S.A., Vevey, Switzerland.


© 2018 Nestlé NEST-14554-1118

\*BOOST Glucose Control® is also available for people with diabetes.

\*\*Take ARGINAID® Powder separately by mixing one packet with 6-8 fl oz water. Two servings per day of ARGINAID® Powder or IMPACT Advanced Recovery® Drink are recommended.



# Nutritional Management of Pressure Injury – Tube Feeding Solutions



**USE UNDER MEDICAL SUPERVISION.**

This document is not intended as a substitute for clinical judgment or existing facility protocols.

Posthauer M et al. *Adv Skin Wound Care* 2015;28(4):175-88.

<http://www.npuap.org/resources/educational-and-clinical-resources/npup-pressure-injury-stages/>

All trademarks are owned by Société des Produits Nestlé S.A., Vevey, Switzerland.

© 2018 Nestlé

\*Mix one packet of ARGINAID® Powder with 4-6 fl oz water and administer by syringe via a feeding tube. Flush tube afterwards with a minimum of 1-2 fl oz water. Two packets per day are recommended. Due to moderate acidity, consult a medical professional before use in a surgical jejunostomy.