


Oral Supplement Selection Guide for Partial and Full Thickness Wounds[¥]



These suggested guidelines are based on clinical references relating to the use of arginine and protein for the nutritional management of wounds, and should not be construed as a substitute for medical advice or existing facility protocols.

[¥] In patients with multiple wounds, the algorithm is directed to the most severe wound.
Reassess wound progress at 4 weeks⁷

*Unintended weight loss = >3kg (or >6 lbs) in past 3 months

USE UNDER MEDICAL SUPERVISION

References:

1. Kaiser MJ et al. *J Nutr Health Aging* 2009;13:782-788.
2. EPUAP, NPIAP & PPPPIA. Prevention and Treatment Pressure Ulcers/Injuries: Quick Reference Guide. Emily Haesler (Ed.). EPUAP/NPIAP/PPPPIA: 2019.
3. Desneves KJ et al. *Clin Nutr* 2005;24:979-987.
4. Huhmann M et al. Case Summary of Data on File 2015.
5. Escott-Stump S. *Nutrition and Diagnosis-Related Care*. 7th ed. Baltimore, MD: Lippincott Williams & Wilkins; 2011.
6. Chow O & Barbul A. *Adv Wound Care* 2014;3(1):46-53.
7. Sheehan P et al. *Diabetes Care*. 2003;26(6):1879-1882.

All trademarks are owned by Société des Produits Nestlé S.A., Vevey, Switzerland. ©2020 Nestlé.
All rights reserved. NEST-14781-0120

